

Grey platelet syndrome

What is it?


- Grey platelet syndrome (GPS) is a rare bleeding disorder which can be diagnosed from childhood to old age.
- GPS is caused by mutations in the *Neurobeachin-like 2 (NBEAL-2)* gene.
- GPS was first described in 1971 by the American haematologist, Dr Giovanni Raccuglia but it was not until 2011 that it shown by three research group to be caused by mutations in *NBEAL-2*.
- There are less than one hundred patients worldwide, and less than ten patients in the UK.
- NBEAL-2 is a protein which is required for the normal assembly of the contents of a particular type of storage granule in platelets known as an α -granule. The α -granules release proteins that support platelet aggregation (clumping) and which also help to repair the injury and prevent infection.
- Mutations in *NBEAL-2* cause a mild reduction in the number of platelets (thrombocytopenia) and a mild defect in platelet function.
- NBEAL-2 is only found in platelets and the platelet mother cell, the megakaryocyte.

Who suffers?

- GPS affects equal numbers of males and females. Two copies of the mutant gene are inherited (one each) from the mother and father (autosomal recessive). It is therefore more common in families where individuals are blood relatives of each other. In general carriers of the condition (who only have one mutated gene) do not experience clinical problems.

What are the symptoms?

- Patients with GPS may experience nose bleeding, easy bruising, bleeding from gums, heavy or prolonged menstrual bleeding (menorrhagia), bleeding after childbirth, abnormal bleeding after surgery or dental work, and gastrointestinal bleeding.



©2016 The Platelet Charity

Helping people with bleeding disorders

www.plateletcharity.co.uk

Registered Charity Number: 1151168

Registered Address : Curo House, Greenbox, Westonhall Road, Stoke Prior, Bromsgrove, B60 4AL.


Diagnosis

- Patients have a low platelet count (thrombocytopenia) and enlarged platelets.
- Platelets appear grey on a blood film when examined. This is one of the first tests performed in a hospital. Other platelet disorders can also have grey platelets.
- Platelets may undergo normal platelet aggregation (clumping) but have reduced α -granule secretion to all stimuli which can be detected by a technique called flow cytometry using an antibody to a protein called P-selectin.
- A definitive diagnosis of GPS is made by genetic sequencing.

Treatment

- Treatment should be led by and discussed with a haematologist with experience in bleeding disorders.
- Treatment is the same as other conditions where there are platelet function defects. General strategies include avoidance of medications that inhibit platelet function further (e.g. aspirin) and compression at sites of injury. Patients may be treated with antifibrinolytics (e.g. tranexamic acid) or platelet transfusion prior to surgery.
- Patients should carry a bleeding disorders / haemorrhagic states card (issued by a Haemophilia Care Centre) to alert others to their condition.
- Genetic counselling may be offered.

The Platelet Charity 2017



©2016 The Platelet Charity

Helping people with bleeding disorders

www.plateletcharity.co.uk

Registered Charity Number: 1151168

Registered Address : Curo House, Greenbox, Westonhall Road, Stoke Prior, Bromsgrove, B60 4AL.